

Counting incompressible surfaces in 3-manifolds

Nathan M. Dunfield
University of Illinois

joint with
Stavros Garoufalidis
Hyam Rubinstein

Throughout: M^3 is a cpt orient irreducible with every closed $F^2 \subset M$ orient (e.g. $H_2(M; \mathbb{F}_2) = 0$).

Closed conn embedded $F^2 \subset M$ is *incompressible* when $F \neq S^2$ and $\pi_1 F \rightarrow \pi_1 M$ is injective; if F is also not parallel into ∂M , it is *essential*.

Goal: Count (closed) essential surfaces in M , up to isotopy.

T^3 : all essential surfaces are tori, infinitely many.


$|\pi_1 M| < \infty$: no essential surfaces.

[Hatcher-Thurston 1985] 2-bridge knot exterior has no ess. surfaces.

M^3 is atoroidal when there are no ess. tori. For atoroidal M , this is always finite:

$$a_M(g) = \#\{\text{genus } g \text{ ess. surf, mod iso}\}$$

For the exterior M of $11n34$:



g	a_M	g	a_M	g	a_M
1	0	7	87	13	602
2	6	8	208	14	1,168
3	9	9	220	15	1,039
4	24	10	366	16	1,498
5	37	11	386	17	1,564
6	86	12	722	18	2,514
					...
				50	56,892
				100	444,038

$$b_M(-n) = \#\left\{ \begin{array}{l} \text{ess. surf with } \chi = -n \\ \text{mod isotopy} \end{array} \right\}$$

For $M = E_{11n34}$, we show

$$b_M(-2n) = \frac{2}{3}n^3 + \frac{9}{4}n^2 + \frac{7}{3}n + \frac{7 + (-1)^n}{8}$$


Thm [DGR] For atoroidal M^3 , the generating function

$$\sum_{n=1}^{\infty} b_M(-2n)x^n = \frac{P(x)}{Q(x)}$$


where $P, Q \in \mathbb{Q}[x]$ and Q is a product of cyclotomics.

Algorithm [DGR] Can find P, Q , and isotopy reps for fixed χ .

Normal surfaces meet each tetrahedra in a standard way:



and correspond to lattice points in a finite polyhedral cone P_T in \mathbb{R}^{7t} where $t = \#T$:



Good: Any essential F can be isotoped to be normal.

Bad: Resulting normal surface is far from unique.

weight: $\text{wt}(F) = \#(F \cap T^1)$

lw-surface: an essential normal surface that is least weight in its isotopy class.

[Tollefson 90s, Oertel 80s]

Every lw-surface lies on a lw-face $C \subset P_T$, one where **every** lattice point in C is a lw-surface. Isotopies between lw-surfaces can be understood.

[Ehrhart 60s] Counts of lattice points in rational polyhedra are quasipolynomial.

Thm [DGR] For atoroidal M^3 , the count $b_M(-n)$ is quasipolynomial.

Moral: Ess. surf. are lattice points in the space $\mathcal{ML}(M)$ of measured laminations [Hatcher '90s].

Cor [DGR] The number of ess. surfaces of $\chi = -2n$ grows like n^{d-1} where $d = \dim(\mathcal{ML}(M))$.

[Kahn-Markovic 2012] For M^3 closed hyperbolic, the number of **immersed** essential genus g surfaces grows like g^{2g} .

What about counting by genus? Which lattice points correspond to connected surfaces? Even for curves on surfaces, this is very subtle [Mirzakhani].

Computed $\mathcal{LW}_T = \bigcup \{C \text{ is a lw-face}\}$ for 59K manifolds. Some 4K with $\dim(\mathcal{LW}_T) > 1$.

For $K13n3838$, \mathcal{LW}_T is conn. with 44 maximal faces, all of dim 5, each with 5–9 vertex rays cor. to 48 distinct surfaces of genus 2–5. Here $b_M(-2n)$ is:

$$\frac{7}{12}n^4 + 3n^3 + \frac{14}{3}n^2 + 3n + \frac{7 + (-1)^n}{8}$$

and $a_M(g)$ starts 12, 34, 110, 216, 532, 708, 1558, 2018, 3462, 4176, 7314, 7876, 13204, 14256, 20778, 23404, 34820, 34832, 52226,...