Math 291 Workshop 6

Please work in groups of two or three. Please explain all answers carefully.

Be sure to acknowledge any and all help you receive with your write-up. As usual, you can *think* with other people but you should not *write* with anybody. Please give your answer in complete English sentences. Helpful figures are always welcome.

**Problem 6.1 (Warm-up).** The constraint  $x^2 + 2y^2 \le 1$  defines a closed and bounded set in  $\mathbb{R}^2$ . Thus, the function f(x,y) = xy attains a maximum value on that set. What is this maximum value? Be sure to analyze carefully and completely any system of equations you solve.

**Problem 6.2 (Warm-up).** Suppose that a and b are fixed positive real numbers. The constraint  $ax^2 + by^2 \le 1$  defines a closed and bounded set in  $\mathbb{R}^2$ . Thus, the function f(x,y) = xy attains a maximum value on that set. What is this maximum value? Explain.

**Problem 6.3.** Let G be the graph of the function z = xy, in  $\mathbb{R}^3$ . Sketch G. For every real number a, let  $P_a = (0,0,a)$ . Now find the minimal distance between  $P_a$  and G. (Hint: it is a bit easier to minimize the square of the distance.) Discuss the geometric meaning of all of the non-minimal solutions found by the Lagrange multiplier method.

**Problem 6.4.** Let G be the graph of the function  $z = x^2 + 2y^2$ . Sketch G in  $\mathbb{R}^3$ . Let  $P_a = (0,0,a)$  and find the minimal distance between  $P_a$  and G. Explain geometrically the various other solutions given by the Lagrange method. In particular, why do different values of a lead to different numbers of critical points?

**Problem 6.5.** Suppose that, in the problem above, G is the graph of  $z = x^2 + y^2$ . Draw a picture. What do you expect to happen in this case? How many critical points will there be when a is large and positive? When a is small and positive?

Now solve the problem using Lagrange. At what value of a does the behavior change? Why? Explain geometrically.

2007/03/01